
What is claimed is:

## 1. A compound of the Formula (I)



wherein

5

10

15

20

25

30

Z is a single bond or a  $C_1$ - $C_4$  hydrocarbon chain containing no more than 1 double or triple bond, optionally substituted with one or more substituents independently selected from the group consisting of  $C_1$ - $C_4$  alkyl;

A is an aromatic ring selected from the group consisting of optionally substituted arylene and optionally substituted heteroarylene, wherein A is not benzimidazole and when Z is a single bond then A is not selected from the group consisting of phenylene and six-membered heteroarylene containing 3 or less than 3 nitrogens;

B is an aromatic ring selected from the group consisting of optionally substituted aryl, optionally substituted arylene, optionally substituted heteroarylene and wherein A and B can not both be phenylene and wherein when Z is a single bond then B is not a bicyclic aryl or bicyclic heteroaryl;

wherein A and B are connected via a carbon-carbon bond;

R<sub>2</sub> is selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylalkenyl, cycloalkylheteroalkyl, heterocycloalkylheteroalkyl, heteroarylheteroalkyl, arylheteroalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkenyloxy, alkynyloxy, cycloalkylkoxy, heterocycloalkyloxy, aryloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, aminoalkyl, acylamino, arylamino, phenoxy, benzyloxy, COOH, COOR4, SH, CONHR4, NHR4, -(CH2)nNHCOR4, NHCOR4. NHCOOR4 NHCONHR4, C(=NOH)R4, NHSOR4 NHSO2R4, -(CH2)n-NR6R7, alkoxycarbonyl.

106

alkylaminocarbonyl, sulfonyl, alkylsulfonyl, alkylsulfinyl, arylsulfonyl, arylsulfinyl, aminosulfonyl, aminosulfinyl,  $SR_4$  and acyl each of which may optionally be substituted, provided that  $R_2$  does not contain the moiety NHCONHCO or NHCONHSO<sub>2</sub>;

R<sub>3</sub> is selected from the group consisting of H, halogen, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyi, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylalkenyl, cycloalkylheteroalkyl. heterocycloalkylheteroalkyl. heteroarylheteroalkyl, arylheteroalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkoxyaryl, alkenyloxy, alkynyloxy, cycloalkylkoxy, heterocycloalkyloxy, aryloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, aminoalkyl, acylamino, arylamino, phenoxy, benzyloxy, COOH, COOR4, SH, CONHR4, NHR4, -(CH2)nNHCOR4, NHCOR4, NHCOOR<sub>4</sub> NHCONHR<sub>4</sub>, C(=NOH)R<sub>4</sub>, NHSOR<sub>4</sub> NHSO<sub>2</sub>R<sub>4</sub>, -(CH<sub>2</sub>)<sub>n</sub>-NR<sub>6</sub>R<sub>7</sub>, alkoxycarbonyl, alkylaminocarbonyl, sulfonyl, alkylsulfonyl, alkylsulfinyl, arylsulfonyl, arylsulfinyl, aminosulfonyl, aminosulfinyl, SR4 and acyl; each of which may optionally be substituted provided that R<sub>3</sub> does not contain the moiety NHCONHCO or NHCONHSO<sub>2</sub>;

or  $R_2$  and  $R_3$  together with portion of ring B may form a non-aromatic ring fused to B;

X and Y are the same or different and are independently selected from the group consisting of H, halogen, -CN, -NO<sub>2</sub>, -CF<sub>3</sub>, -OCF<sub>3</sub>, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl. heterocycloalkenyl, heteroaryl, hydroxy, hydroxyalkyl, alkoxyalkyl, alkoxyaryl, alkoxyheteroaryl, alkenyloxy, cycloalkyloxy, alkynyloxy, cycloalkenyloxy, heterocycloalkyloxy, heterocycloalkenyloxy, aryloxy, heteroaryloxy. arylalkyl. heteroarylalkyl, arylalkyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, aminoalkyl, alkoxyalky, -COOH, -C(O)OR4, -COR4, -SH, -SR4, -OR4, acvl and -NR8R6 each of which may be optionally substituted;

30

5

10

15

20

25

each R<sub>4</sub> is independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl and acyl each of which may be optionally substituted;

35

each R<sub>6</sub> and R<sub>7</sub> is independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, cycloalkyl, heteroacyl,

cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl and acyl; each of which may be optionally substituted;

each R<sub>8</sub> and R<sub>9</sub> is independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl and acyl; each of which may be optionally substituted;

n is an integer from 0 to 6,

WO 2005/040161


10

5

m is an integer from 0 to 4;

or a pharmaceutically acceptable salt or prodrug thereof.

## 15 2. A compound according to claim 1 having the Formula (la)



wherein

25

30

Z is a single bond or a  $C_1$ - $C_4$  hydrocarbon chain which may contain 0 to 1 double or triple bonds, unsubstituted or substituted with one or more substituents independently selected from the group consisting of  $C_1$ - $C_4$  alkyl;

A is an aromatic ring selected from the group consisting of optionally substituted arylene and optionally substituted heteroarylene, wherein A is not benzimidazole and when Z is a single bond then A is not selected from the group consisting of phenylene and six-membered heteroarylene containing 3 or less than 3 nitrogens;

B is an aromatic ring selected from the group consisting of optionally substituted aryl, optionally substituted arylene, optionally substituted heteroaryl and optionally

108

substituted heteroarylene and wherein A and B can not both be phenylene and wherein when Z is a sing le bond then B is not a bicyclic aryl or bicyclic heteroaryl;

wherein A and B are connected via a carbon-carbon bond;

5

10

15

R<sub>2</sub> is selected from C<sub>1</sub>-C<sub>10</sub> alkyl, alkenyl, heteroalkyl, haloalkyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, C<sub>4</sub>-C<sub>9</sub> heterocycloalkylalkyl, cycloalkylalkyl cyclopropylmethyl), arylalkyl (e.g. benzyl), heteroarylalkyl (e.g. pyridylmethyl), hydroxyl, hydroxyalkyl, alkoxy, amino, alkylamino, aminoalkyl, acylamino, phenoxy, alkoxyalkyl, benzyloxy, alkylsulfonyl, arylsulfonyl, aminosulfonyl, -C(O)OR4, -C(O)OH, -SH, -CONHR4, -NHCONHR<sub>4</sub>, C(=NOH)R<sub>4</sub>, -C(O)C(O)OR4, C(O)CONHR4, CON(R<sub>5</sub>)OR<sub>4</sub>, COCON(R4)OR4, NHCOR4, and acyl; each of the above is unsubstituted or optionally substituted with one or more substituents independently selected from the group consisting of: halogen; =0; =S; -CN; and -NO<sub>2</sub>; and alkyl, alkenyl, heteroalkyl, haloalkyl, alkynyl, aryl, cycloaikyl, heterocycloaikyl, heteroaryl, hydroxyl, hydroxyalkyl, alkoxy, alkylamino, amimoalkyl, acylamino, phenoxy, alkoxyalkyl, benzyloxy, alkylsulfonyl, arylsulfonyl, aminosulfonyl, -C(O)OR5, -C(O)OH, -SH, -C(O)C(O)OR5, C(O)CONHR5,  $CON(R_5)OR_5$ ,  $COCON(R_5)OR_5$ ,  $NHCOR_5$ , and acyl; wherein  $R_2$  does not contain the moiety NHCONHCO or NHCONHSO2;

20

25

30

R<sub>3</sub> is selected from H, C<sub>1</sub>-C<sub>10</sub> alkyl, alkenyl, heteroalkyl, haloalkyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, C<sub>4</sub>-C<sub>9</sub> heterocycloalkylalkyl, cycloalkylalkyl (e.g., cyclopropylmethy I), arylalkyl (e.g. benzyl), heteroarylalkyl (e.g. pyridylmethyl), hydroxyl, hydroxyalkyl, alkoxy, amino, alkylamino, aminoalkyl, acylamino, phenoxy, alkoxyalkyl, benzyloxy, alkylsulfonyl, arylsulfonyl, aminosulfonyl, -C(O)OR4, -C(O)OH, -SH, -CONHR4, -NHCONHR₄,  $C(=NOH)R_4$ -C(O)C(O)OR<sub>4</sub>, C(O)CONHR4, COCON(R<sub>4</sub>)OR<sub>4</sub>, NHCOR<sub>4</sub>, and acyl; each of the above is unsubstituted or optionally substituted with one or more substituents independently selected from the group consisting of: halogen; =O; =S; -CN; and -NO2; and alkyl, alkenyl, heteroalkyl, haloalkyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, hydroxyl, hydroxyalkyl, alkoxy. alkylamino, aminoalkyl, acylamino, phenoxy, alkoxyalkyl, benzyloxy, alkylsulfonyl, arylsulfonyl, aminosulfonyl, -C(0)OR<sub>5</sub>, -C(0)OH, -SH, -C(0)C(0)OR<sub>5</sub>, C(0)CONHR<sub>5</sub>.  $CON(R_5)OR_5$ ,  $COCON(R_5)OR_5$ ,  $NHCOR_5$ , and acyl; wherein  $R_3$  does not contain the moiety NHCONHCO or NHCONHSO2;

35

B;

or R<sub>2</sub> and R<sub>3</sub> together with portion of ring B may form a non-aromatic ring fused to

X and Y are the same or different and independently selected from the group consisting of: H, halo,  $C_1$  - $C_4$  alkyl, such as  $CH_3$  and  $CF_3$ ,  $NO_2$ ,  $OR_4$ ,  $SR_4$ ,  $C(O)R_5$ , CN, and  $NR_8$   $R_9$ ;

5

R<sub>4</sub> is selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, heteroalkyl, aryl, heteroaryl, acyl;

R<sub>5</sub> is selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl;

10


 $R_8$  and  $R_9$  are the same or different and independently selected from the group consisting of H,  $C_1$ - $C_6$  alkyl,  $C_4$ - $C_9$  cycloalkyl,  $C_4$ - $C_9$  heterocycloalkyl, aryl, heteroaryl, arylalkyl, and heteroarylalkyl;

m is an integer from 0 to 4;

15

or a pharmaceutically acceptable salt or prodrug thereof.

3. A compound according to claim 1 or 2 having the Formula (lb)



Formula (lb)

20 wherein

Z is a single bond or a  $C_1$ - $C_4$  hydrocarbon chain which may contain 0 to 1 double bond or triple bond, unsubstituted or substituted with one or more substituents independently selected from the group consisting of  $C_1$ - $C_4$  alkyl;

25

30

A is an optionally substituted five-membered heteroarylene:

B is an aromatic ring which is selected from the group consisting of optionally substituted aryl, optionally substituted arylene or optionally substituted heteroaryl or optionally substituted heteroarylene; wherein when Z is a single bond then B is not a bicyclic aryl or bicyclic heteroaryl;

10

15

20

25

35

## wherein A and B are connected via a carbon-carbon bond;

R<sub>2</sub> is selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, haloalkyl. haloalkenyl. heteroalkyi, cycloalkyl, cycloalkenyl, heterocycloalkyl. heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylalkenyl, cycloalkylheteroalkyl, heterocycloalkylheteroalkyl. heteroarylheteroalkyl, arylheteroalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkoxyaryl, alkenyloxy, alkynyloxy, cycloalkylkoxy, heterocycloalkyloxy, aryloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, aminoalkyl, acylamino, arylamino, phenoxy, benzyloxy, COOH, COOR<sub>4</sub>, SH, CONHR<sub>4</sub>, NHR<sub>4</sub>, -(CH<sub>2</sub>)<sub>n</sub>NHCOR<sub>4</sub>, NHCOR<sub>4</sub>, NHCOOR<sub>4</sub> NHCONHR<sub>4</sub>, C(=NOH)R<sub>4</sub>, NHSOR<sub>4</sub> NHSO<sub>2</sub>R<sub>4</sub>, -(CH<sub>2</sub>)<sub>n</sub>-NR<sub>6</sub>R<sub>7</sub>, alkoxycarbonyl, alkylaminocarbonyl, sulfonyl, alkylsulfonyl, alkylsulfinyl, arylsulfonyl, arylsulfinyl, aminosulfonyl, aminosulfinyl, SR4 and acyl; each of which may optionally be substituted, wherein R<sub>2</sub> does not contain the moiety NHCONHCO or NHCONHSO<sub>2</sub>;

R<sub>3</sub> is selected from the group consisting of H, halogen, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, cycloalkyl, heteroalkyl, cycloalkenyl. heterocycloalkyl. heterocycloalkenyi, aryi, heteroaryi, cycloalkylalkyi, heterocycloalkylalkyi, aryialkyi, heteroarylalkyl. arylalkenyl, cycloalkylheteroalkyl. heterocycloalkylheteroalkyl, heteroarylheteroalkyl, arylheteroalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkenyloxy, alkynyloxy, cycloalkylkoxy, heterocycloalkyloxy, aryloxy, alkoxvarvl. heteroaryloxy, arylalkyloxy, amino, alkylamino, aminoalkyl, acylamino, arylamino, phenoxy, benzyloxy, COOH, COOR4, SH, CONHR4, NHR4, -(CH2)nNHCOR4, NHCOR4, NHCOOR4 NHCONHR4, C(=NOH)R4, NHSOR4 NHSO<sub>2</sub>R4, -(CH<sub>2</sub>)<sub>n</sub>NR<sub>6</sub>R<sub>7,</sub> alkoxycarbonyl, alkylaminocarbonyl, sulfonyl, alkylsulfonyl, alkylsulfinyl, arylsulfonyl, arylsulfinyl, aminosulfonyl, aminosulfinyl, SR4 and acyl; each of which may optionally be substituted wherein R<sub>3</sub> does not contain the moiety NHCONHCO or NHCONHSO<sub>2</sub>;

or  $R_2$  and  $R_3$  together with portion of ring B may form a non-aromatic ring fused to B;

X and Y are the same or different and are independently selected from the group consisting of H, halo, C<sub>1</sub> -C<sub>4</sub> alkyl, such as CH<sub>3</sub> and CF<sub>3</sub>, NO<sub>2</sub>, OR<sub>4</sub>, SR<sub>4</sub>, C(O)R<sub>5</sub>, CN, and NR<sub>8</sub> R<sub>9</sub>;

R<sub>4</sub> is selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, heteroalkyl, aryl, heteroaryl, acyl;

10

15

20

25

30

R<sub>5</sub> is selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl;

each R<sub>6</sub> and R<sub>7</sub> is independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl and acyl each of which may be optionally substituted;


 $R_8$  and  $R_9$  are the same or different and are independently selected from the group consisting of H,  $C_1$ – $C_6$  alkyl,  $C_4$ - $C_9$  cycloalkyl,  $C_4$ - $C_9$  heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl;

n is an integer from 0 to 6;

m is an integer from 0 to 4;

or a pharmaceutically acceptable salt or prodrug thereof.

4. A compound according to claim 1 or 2 having the compound of Formula (lc):



wherein

Z is a single bond or a  $C_1$ - $C_4$  hydrocarbon chain which may contain 0 to 1 double bond or triple bond, unsubstituted or substituted with one or more substituents independently selected from the group consisting of  $C_1$ - $C_4$  alkyl;

A is a six-membered aromatic ring which is selected from the group consisting of optionally substituted arylene or optionally substituted heteroarylene and when Z is a single bond then A is not selected from the group consisting of phenylene and six-membered heteroarylene containing 3 or less than 3 nitrogens;

112

B is an aromatic ring and is attached to the 3rd or 4<sup>th</sup> position relative to Z of ring A selected from the group consisting of optionally substituted aryl, optionally substituted arylene, optionally substituted heteroaryl and optionally substituted heteroarylene and wherein A and B can not both be phenylene;

5

10

15

wherein A and B are connected via a carbon-carbon bond;

R<sub>2</sub> is selected from the group consisting of halogen, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, heteroalkyl, cycloalkyl. cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylalkenyl. cycloalkylheteroalkyl. heterocycloalkylheteroalkyl, heteroarylheteroalkyl, arylheteroalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkoxyaryi, alkenyloxy, alkynyloxy, cycloalkylkoxy, heterocycloalkyloxy, aryloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, aminoalkyl, acylamino, arylamino, phenoxy, benzyloxy, COOH, COOR4, SH, CONHR4, NHR4, -(CH2)nNHCOR4, NHCOR4, NHCOOR<sub>4</sub> NHCONHR<sub>4</sub>, C(=NOH)R<sub>4</sub>, NHSOR<sub>4</sub> NHSO<sub>2</sub>R<sub>4</sub>, -(CH<sub>2</sub>)<sub>n</sub>-NR<sub>6</sub>R<sub>7</sub>, alkoxycarbonyl, alkylaminocarbonyl, sulfonyl, alkylsulfonyl, alkylsulfinyl, arylsulfonyl, arylsulfinyl, aminosulfonyl, aminosulfinyl, SR4 and acyl; each of which may optionally be substituted, wherein R<sub>2</sub> does not contain the moiety NHCONHCO or NHCONHSO<sub>2</sub>:

20

25

R<sub>3</sub> is selected from the group consisting of H, halogen, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl. heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylalkenyl, cycloalkylheteroalkyl, heterocycloalkylheteroalkyl, heteroarylheteroalkyl, arylheteroalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkoxyaryi, alkenyloxy, alkynyloxy, cycloalkylkoxy, heterocycloalkyloxy, heteroaryloxy, arylalkyloxy, amino, alkylamino, aminoalkyl, acylamino, arylamino, phenoxy, benzyloxy, COOH, COOR4, SH, CONHR4, NHR4, -(CH2)nNHCOR4, NHCOR4, NHCOOR<sub>4</sub> NHCONHR<sub>4</sub>, C(=NOH)R<sub>4</sub>, NHSOR<sub>4</sub> NHSO<sub>2</sub>R<sub>4</sub>, -(CH<sub>2</sub>)<sub>n</sub>NR<sub>6</sub>R<sub>7</sub>, alkoxycarbonyl, sulfonyl, alkylsulfonyl, arylsulfonyl, arylsulfonyl, arylsulfonyl, alkylaminocarbonyl. aminosulfonyl, aminosulfinyl, SR4 and acyl; each of which may optionally be substituted wherein R<sub>3</sub> does not contain the moiety NHCONHCO or NHCONHSO<sub>2</sub>;

X and Y are the same or different and independently selected from H, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, such as CH<sub>3</sub> and CF<sub>3</sub>, NO<sub>2</sub>, OR<sub>4</sub>, SR<sub>4</sub>, C(O)R<sub>5</sub>, CN, and NR<sub>8</sub> R<sub>9</sub>;

35

30

R<sub>4</sub> is selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, heteroalkyl, aryl, heteroaryl, acyl;

R<sub>5</sub> is selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl;

each R<sub>6</sub> and R<sub>7</sub> is independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl and acyl each of which may be optionally substituted;

 $R_8$  and  $R_9$  are the same or different and independently selected from H,  $C_1$ - $C_6$  alkyl,  $C_4$ - $C_9$  cycloalkyl,  $C_4$ - $C_9$  heterocycloalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl;


10

5

n is an integer from 0 to 6;

m is an integer from 0 to 4;

- or a pharmaceutically acceptable salt or prodrug thereof.
  - 5. A compound according to claim 1 having the Formula (Id):



$$\begin{cases} W_{1} & W_{2} & W_{3} \\ W_{2} & W_{3} \\ W_{2} & W_{3} \\ W_{3} & W_{3} \\ W_{4} & W_{3} \\ W_{5} & W_{3} \\ W_{5} & W_{3} \\ W_{5} & W_{3} \\ W_{5} & W_{5} \\ W_$$

wherein W<sub>1</sub> is selected from the group consisting of O, S and NH;

W<sub>2</sub> and W<sub>3</sub> are independently selected from the group consisting of N, CX and CY;

p is an integer from 0 to 3,


5

15

20

B is a 5-membered heteroarylene,

- wherein Z, X, Y, R<sub>2</sub> and R<sub>3</sub> are as described in claim 1, or a pharmaceutically acceptable salt or prodrug thereof.
  - 6. A compound according to claim 1 having the Formula (le):



Formula (le)

wherein B is a 5-membered heteroarylene, p is an integer from 0 to 3 and X, Y,  $R_2$  and  $R_3$  are the same as in claim 1.

7. A compound according to claim 1 having the Formula (If)

15

20

$$R_2$$
 $B$ 
 $(CH_2)$ 
 $C$ 
 $OH$ 
 $N$ 
 $N$ 
 $H$ 

Formula (If)

wherein B is a 5-membered heteroarylene, p is an integer from 0 to 3 and X, Y,  $R_2$  and  $R_3$  are the same as in claim 1.

8. A compound according to claim 1 of the Formula (lg):

Formula (Ig)


wherein q is an integer from 0 to 4 and X, Y, R<sub>2</sub> and R<sub>3</sub> are the same as in claim 1.

9. A compound according to claim 1 of the Formula (lh):

Formula (lh)

wherein q is an integer from 0 to 4 and X, Y, R<sub>2</sub> and R<sub>3</sub> are the same as in claim 1.

10. A compound according to claim 1 of the Formula (li):




Formula (li)

X, Y,  $R_2$  and  $R_3$  are the same as in claim 1.

5

## 11. A compound according to claim 1 of the Formula (Ij):



10

r is an integer from 0 to 4 and X, Y,  $R_2$  and  $R_3$  are the same as in claim 1.

12. A compound according to claim 1 of the Formula (lk):

r is an integer from 0 to 4 and X, Y, R<sub>2</sub> and R<sub>3</sub> are the same as in claim 1.

5

10

15

20

25

13. A compound according to any one of claims 1 to 3 wherein A is an optionally substituted 5-membered heteroarylene ring.

- 14. A compound according to any one of claims 1 to 3 or 5 wherein A is an optionally substituted 5-membered heteroarylene ring selected from the group consisting of 2.5furanylene; 2,4-furanylene; 2,3-furanylene; 3,4-furanylene; 2,5-thiophenylene; 2,4thiophenylene, 2,3-thiophenylene; 3,4-thiophenylene; 1,2-pyrrolylene; 1,3-pyrrolylene; 1,4pyrrolylene; 1,5-pyrrolylene; 2,3-pyrrolylene; 2,4-pyrrolylene; 3,4pyrrolylene; 2,5-oxazolylene; 2,4-oxazolylene; 4,5-oxazolylene, 2,5-thiazolylene; 2,4thiazolylene; 4,5-thiazolylene 1,2-imidazolylene; 1,4-imidazolylene; 1,5-imidazolylene; 2,4imidazolylene; 2,5- imidazolylene; 4,5-imidazolylene 1,3-pyrazolylene; 1,4-pyrazolylene; 1,5-pyrazolylene; 3,4-pyrazolylene; 3,5-pyrazolylene; 4,5-pyrazolylene; 3,4-isoxazolylene; 3,5-isoxazolylene; 4,5-isoxazolylene; 3,4-isothiazolylene; 3,5-isothiazolylene; 4,5isothiazolylene; 4,5-(1,2,3-oxadiazoly)-ene; 3,5,-(1,2,4-oxadiazolyi)ene; 1,4-(1,2,3triazolyl)ene; 1,5-(1,2,3-triazolyl)ene; 4,5-(1,2,3-triazolyl)ene; 1,3-(1,2,4-triazolyl)ene; 1,5-(1,2,4-triazolyl)ene; 3,5-(1,2,4-triazolyl)ene; 3,5-(1,2,4-thiadiazolyl)ene; thiadiazolyl)ene, and 1,5-tetrazolylene.
- 15. A compound according to any one of claims 1 to 3 or 5 wherein A is an optionally substituted 5-membered heteroarylene selected from the group consisting of 2,5-thiophenylene; 3,5-isoxazolylene; 3,5-pyrazolylene; 2,5-oxazolylene; 3,5-pyrazolylene; 2,5-furanylene and 2,4-thiophenylene.
- 16. A compound according to any one of claims 1 to 3 or 5 wherein B is attached to the 3<sup>rd</sup> or 4<sup>th</sup> position relative to Z of Ring A.

- 17. A compound according to any one of claims 1, 2 or 4 wherein A is an optionally substituted phenylene or an optionally substituted 6-membered heteroarylene.
- 5 18. A compound according to any one of claims 1 to 7 or 13 to 17 wherein B is an optionally substituted 5-membered heteroarylene.

10

15

20

25

- A compound according to any one of claims 1 to 7 or 13 to 18 wherein B is an 19. optionally substituted 5-membered heteroarylene ring selected from the group consisting of 2,5-furanylene; 2,4-furanylene; 2,3-furanylene; 3,4-furanylene; 2,5-thiophenylene; 2,4thiophenylene, 2,3-thiophenylene; 3,4-thiophenylene; 1,2-pyrrolylene; 1,3-pyrrolylene; 1,4pyrrolylene; 1,5-pyrrolylene; 2,3-pyrrolylene; 2,4-pyrrolylene; 2,5-pyrrolylene; 3,4pyrrolylene; 2,5-oxazolylene; 2,4-oxazolylene; 4,5-oxazolylene, 2,5-thiazolylene; 2,4thiazolylene; 4,5-thiazolylene 1,2-imidazolylene; 1,4-imidazolylene; 1,5-imidazolylene; 2,4imidazolylene; 2,5- imidazolylene; 4,5-imidazolylene 1,3-pyrazolylene; 1,4-pyrazolylene; 1,5-pyrazolylene; 3,4-pyrazolylene; 3,5-pyrazolylene; 4,5-pyrazolylene; 3,4-isoxazolylene; 3,5-isoxazolylene; 4,5-isoxazolylene; 3,4-isothiazolylene; 3,5-isothiazolylene; 4,5isothiazolylene; 4,5-(1,2,3-oxadiazoly)-ene; 3,5,-(1,2,4-oxadiazolyl)ene: 1,4-(1,2,3triazolyl)ene; 1,5-(1,2,3-triazolyl)ene; 4,5-(1,2,3-triazolyl)ene; 1,3-(1,2,4-triazolyl)ene; 1,5-(1,2,4-triazolyl)ene; 3,5-(1,2,4-triazolyl)ene; 3,5-(1,2,4-thiadiazolyl)ene; 2,5-(1,3,4thiadiazolyl)ene, and 1,5-tetrazolylene.
- 20. A compound according to any one of claims 1 to 7 or 13 to 19 wherein B is an optionally substituted 5-membered heteroarylene selected from the group consisting of 2,4-thiazolylene; 4,2-thiazolylene; 1,3-phenylene; 2,5-thiophenylene and 1,4-phenylene.
- 21. A compound according to any one of claims 1 to 5 wherein Z is a single bond.
- 22. A compound according to any one of claims 1 to 5 wherein Z is CH<sub>2</sub>.
- 23. A compound according to any one of claims 1 to 5 wherein Z is CH=CH.
- 24. A compound according to any one of claims 1 to 23 wherein X and Y are both H.
- 25. A compound according to any one of the preceding claims wherein R<sub>2</sub> is independently iselected from the group consisting of: -NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>n</sub>NHCOR<sub>4</sub>, -NHSO<sub>2</sub>R<sub>4</sub>, -NR<sub>4</sub>, -(CH<sub>2</sub>)<sub>n</sub>NR<sub>6</sub>R<sub>7</sub>-, arylalkyl and heteroarylalkyl, each of which may be optionally

10

15

20

25

substituted wherein n is an integer from 0 to 6, and  $R_4$ ,  $R_6$  and  $R_7$  are as described in claim 1.

- 26. A compound according to any one of the preceding claims wherein  $R_2$  is selected from the group consisting of  $R_6R_7N$ -(CH<sub>2</sub>)<sub>n</sub>- wherein n is an integer from 1 to 3.
- 27. A compound according to claim 26 wherein  $R_6$  and  $R_7$  are independently selected from the group consisting of:
- H, cyclopropyl, 2-(4-Hydroxy-3,5-dimethoxy-phenyl)-ethyl, 3-Pyrrolidin-1-yl-propyl, 2-Morpholin-4-yl-ethyl, 3-Morpholin-4-yl-propyl, 2-Dimethylamino-ethyl. 4-[4-(2,3-Dimethylphenyl)-piperazin-1-ylmethyl, 3-lmidazol-1-yl-propyl, 3-phenyl-propyl, (2-Hydroxy-ethyl)phenethyl, 2-Hydroxy-ethyl-2-(1H-indol-3-yl)-ethyl, (2-Morpholin-4-yl-ethyl)-phenethyl, 2-(2-methyl-1H-indol-3-yl)-ethyl, 2-(1H-indol-3-yl)-ethyl, pyridin-3-ylmethyl, 3-hydroxy-propyl, 2-pyridin-2-yl-ethyl, 2-pyridin-3-yl-ethyl, pyridin-3-ylmethyl, 2-pyridin-4-yl-ethyl, benzyl, 3phenyl-propyl, 2-phenoxy-ethyl, morpholin-4-yl, pyridin-2-yl, phenethyl, 2-(4-bromophenyl)-ethyl, 2-(4-fluoro-phenyl)-ethyl, 3-imidazol-1-yl-propyl, 2-(1H-imidazol-4-yl)-ethyl, 1H-Benzoimidazol-2-ylmethyl, 2-piperidin-1-yl-ethyl, 2-pyrrolidin-1-yl-ethyl, 2-cyclohex-1enyl-ethyl, 2-ethyl-hexyl, 2-thiophen-2-yl-ethyl, 3,3-diphenyl-propyl, 2-biphenyl-4-yl-ethyl, -(4-phenoxy-phenyl, 2-(3-phenoxy-phenyl)-ethyl, 2-(2,3-dimethoxy-phenyl, 2-(2,4-dichlorophenyl)-ethyl, cyclohexylmethyl, hexyl, isobutyl, 3-isopropoxy-propyl, 2-phenoxy-ethyl, 2isopropoxy-ethyl, 3-methoxy-benzyl, 4-[1,2,3]thiadiazol-4-yl-benzyl, 2,4-dichloro-benzyl, 2-(2-methoxy-phenyl)-ethyl, 2-(3-fluoro-phenyl)-ethyl, 2-(2-fluoro-phenyl)-ethyl, 2,2-diphenylethyl, 2-(4-methoxy-phenyl)-ethyl, 2-(3-chloro-phenyl)-ethyl, 4-phenyl-butyl, 3-phenylpropyl, 3,3-diphenyl-propyl, 3-(4-methyl-piperazin-1-yl, 3-morpholin-4-yl-propyl, 3-(2-oxopyrrolidin-1-yl)-propyl, 3-pyrrolidin-1-yl-propyl, tetrahydro-furan-2-ylmethyl, 2-diethylaminoethyl, 2-dimethylamino-ethyl.
- 28. A compound according to claim 1 wherein the compound is selected from the group consisting of

N-Hydroxy-2-[5-(2-phenylacetylamino-thiazol-4-yl)-thiophen-2-yl]-acetamide,

5-(2-Phenylacetylamino-thiazol-4-yl)-isoxazole-3-carboxylic acid hydroxyamide,

PCT/SG2004/000354

120

5-(3-Benzoylamino-phenyl)-1H-pyrazole-3-carboxylic acid hydroxyamide,

2-[5-(2-Benzenesulfonylamino-thiazol-4-yl)-thiophen-2-yl]-N-hydroxy-acetamide,

**{**5-[4-(5-Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-ylcarbamoyl]-benzofuran-2-ylmethyl}-carbamic acid tert-butyl ester,

2-[5-(2-Amino-thiazol-4-yl)-thiophen-2-yl]-N-hydroxy-acetamide,

4-Methyl-2-piperidin-3-yl-thiazole-5-carboxylic acid [4-(5-hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-yl]-amide,

3-{5-[4-(5-Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-ylcarbamoyl]-4-methyl-thiazol-2-yl}-piperidine-1-carboxylic acid tert-butyl ester,

N-[4-(5-2Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-yl]-4-piperazin-1-yl-benzamide,

4-{4-[4-(5-Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-ylcarbamoyl]-phenyl}piperazine-1-carboxylic acid tert-butyl ester,

2-Phenyl-1-piperidin-4-ylmethyl-1H-pyrrole-3-carboxylic acid [4-(5-hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-yl]-amide,

4-{3-[4-(5-Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-ylcarbamoyl]-2-phenyl-pyrrol-1-ylmethyl}-piperidine-1-carboxylic acid tert-butyl ester.

4-{5-[4-(5-Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-ylcarbamoyl]-4-phenyl-thiazol-2yl}-piperidine-1-carboxylic acid tert-butyl ester,

3-[4-(2-Amino-thiazol-4-yl)-phenyl]-N-hydroxy-acrylamide,

3-[4-(2-Acetylamino-thiazol-4-yl)-phenyl]-N-hydroxy-acrylamide,

5-(2-Benzylamino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-(2-Amino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-(3,4-Dimethoxy-phenyl)-oxazole-2-carboxylic acid hydroxyamide,

5-(2-Benzoylamino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-(2-Phenylacetylamino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-(2-Benzenesulfonylamino-thiazol-4-yl)thiophene-2-carboxylic acid hydroxyamide,

5-[2-(2-Phenoxy-acetylamino)-thlazol-4-yl]-thiophene-2-carboxylic acid hydroxyamlde,

PCT/SG2004/000354

WO 2005/040161

122

2-[5-(2-Acetylamino-thiazol-4-yl)-thiophen-2-yl]-N-hydroxy-acetamide,

5-[4-(Phenethylamino-methyl)-phenyl]thiophene-2-carboxylic acid hydroxyamide,

5-[3-(Phenethylamino-methyl)-phenyl]thiophene-2-carboxylic acid hydroxyamide,

5-[2-(3-Phenyl-propylamino)-thiazol-4-yl]-thiophene-2-carboxylic acid hydroxyamide,

5-{4-[(2-Pyridin-2-yl-ethylamino)-methyl]-phenyl}-thiophene-2-carboxylic acid hydroxyamide,

5-{3-[(2-Pyridin-2-yl-ethylamino)-methyl]-phenyl}-thiophene-2-carboxylic acid hydroxyamide,

5-(4-{[2-(1H-Indol-3-yl)-ethylamino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-(3-{[2-(1H-Indol-3-yl)-ethylamino]-methyl}phenyl)thiophene-2-carboxylic acid hydroxyamide,

5-(3-Benzenesulfonylamino-phenyl)-1H-pyrazole-3-carboxylic acid hydroxyamide,

5-(3-Phenylacetylamino-phenyl)-1H-pyrazole-3-carboxylic acid hydroxyamide,

123

2-[5-(3,4-Dimethoxy-phenyl)-oxazol-2-yl]-N-hydroxy-acetamide

3-[5-(3-Chloro-phenyl)-furan-2-yl]-N-hydroxy-acrylamide,

N-Hydroxy-3-{4-[2-(3-phenyl-propyl)-oxazol-5-yl]-phenyl}-acrylamide,

5-[4-({(2-Hydroxy-ethyl)-[2-(1H-indol-3-yl)-ethyl]-amino}-methyl)-phenyl]-thiophene-2-carboxylic acid hydroxyamide,

5-(4-{[(2-Hydroxy-ethyl)-phenethyl-amino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-[3-({(2-Hydroxy-ethyl)-[2-(1H-indol-3-yl)-ethyl]-amino}-methyl)-phenyl]-thiophene-2-carboxylic acid hydroxyamide,

5-(3-{[(2-Morpholin-4-yl-ethyl)-phenethyl-amino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-(3-{[(2-Hydroxy-ethyl)-phenethyl-amino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-{4-[(Pyridin-4-ylmethyl)-amino]-phenyl}-thiophene-2-carboxylic acid hydroxyamide,

HO-NH HO-NH

5-(4-Benzylamino-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-(4-{[(2-Morpholin-4-yl-ethyl)-phenethyl-amino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-{4-[(2-Pyridin-3-yl-ethylamino)-methyl]phenyl}-thiophene-2-carboxylic acid hydroxyamide,

5-{3-[(2-Pyridin-3-yl-ethylamino)-methyl]-phenyl}-thiophene-2-carboxylic acid hydroxyamide,

5-(3-{[(2-Hydroxy-ethyl)-(2-pyridin-3-yl-ethyl)-amino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-[4-(Phenethylamino-methyl)-phenyl]-furan-2-carboxylic acid hydroxyamide,

5-(4-{[2-(1H-Indol-3-yl)-ethylamino]-methyl}phenyl)-furan-2-carboxylic acid hydroxyamide,

5-[3-(Benzylamino-methyl)-phenyl]-furan-2-carboxylic acid hydroxyamide,

5-[3-(Phenethylamino-methyl)-phenyl]-furan-2-carboxylic acid hydroxyamide,

5-{3-[(3-Phenyl-propylamino)-methyl]-phenyl}-furan-2-carboxylic acid hydroxyamide,

4-[4-(Phenethylamino-methyl)-phenyl]-thiophene-2-carboxylic acid hydroxyamide,

4-(4-{[2-(1H-Indol-3-yl)-ethylamino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

or a pharmaceutically acceptable salt or prodrug thereof.

29. A compound according to claim 1 wherein the compound is selected from the group consisting of

N-Hydroxy-2-[5-(2-phenylacetylamino-thiazol-4-yl)-thiophen-2-yl]-acetamide,

5-(2-Phenylacetylamino-thiazol-4-yl)-isoxazole-3-carboxylic acid hydroxyamide,

5-(3-Benzoylamino-phenyl)-1H-pyrazole-3-carboxylic acid hydroxyamide,

O S NH HO NH HO O NH OH OH

2-[5-(2-Benzenesulfonylamino-thiazol-4-yl)-thiophen-2-yl]-N-hydroxy-acetamide,

N-[4-(5-2Hydroxycarbamoylmethyl-thiophen-2-yl)-thiazol-2-yl]-4-piperazin-1-yl-benzamide,

3-[4-(2-Amino-thiazol-4-yl)-phenyl]-N-hydroxy-acrylamide,

PCT/SG2004/000354

HO-<sup>MH</sup>

126

3-[4-(2-Acetylamino-thiazol-4-yl)-phenyl]-N-hydroxy-acrylamide,

5-(2-Benzylamino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-(2-Phenylacetylamino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-(2-Benzenesulfonylamino-thiazol-4-yl)-thiophene-2-carboxylic acid hydroxyamide,

5-[2-(2-Phenoxy-acetylamino)-thiazol-4-yl]-thiophene-

2-carboxylic acid hydroxyamide,

5-[4-(Phenethylamino-methyl)-phenyl]thiophene-2carboxylic acid hydroxyamide,

5-{4-[(2-Pyridin-2-yl-ethylamino)-methyl]-phenyl}-thiophene
-2-carboxylic acid hydroxyamide,

5-(4-Benzylamino-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-(4-{[(2-Morpholin-4-yl-ethyl)-phenethyl-amino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide,

5-{3-[(2-Pyridin-3-yl-ethylamino)-methyl]phenyl}-thiophene-2-carboxylic acid hydroxyamide,

PCT/SG2004/000354

WO 2005/040161

20

127

5-[4-(Phenethylamino-methyl)-phenyl]-furan-2-carboxylic acid hydroxyamide,

5-[3-(Phenethylamino-methyl)-phenyl]-furan-2-carboxylic acid hydroxyamide,

4-[4-(Phenethylamino-methyl)-phenyl]thiophene-2-carboxylic acid hydroxyamide,

4-(4-{[2-(1H-Indol-3-yl)-ethylamino]-methyl}-phenyl)-thiophene-2-carboxylic acid hydroxyamide

or a pharmaceutically acceptable sait or prodrug thereof.

- 30. A compound according to any one of claims 1 to 29 wherein when Z is a single bond then A is not 2,5-thiophenylene.
  - 31. A compound according to any one of claims 1 to 29 wherein when A is phenylene then B is not a 5-membered heteroaryl or 5-membered heteroarylene.
- 32. A compound according to any one of claims 1 to 29 wherein B is not a bicyclic heteroaryl or bicyclic heteroarylene having 9 ring atoms or a hetrocycloalkyl substituted heteroarylene.
- 33. A compound according to any one of claims 1 to 29 wherein when A is a benzimidazole ring, B is not connected to the position 2 of benzimidazole ring.
  - 34. A compound according to any one of claims 1 to 33 wherein the optional substituents are independently selected from the group consisting of H, halogen, =O, =S, -CN, -NO<sub>2</sub>, -CF<sub>3</sub>, -OCF<sub>3</sub>, alkyl, alkenyl, alkynyl, haloalkyl, haloalkynyl, heteroalkyl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl, heteroaryl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkoxyaryl, alkoxyheteroaryl, alkenyloxy, alkynyloxy, cycloalkyloxy, cycloalkenyloxy, heterocycloalkyloxy, heterocycloalkylalkyl, heterocycloalkylalkyl, heterocycloalkylalkyl,

10

20

25

30

arylalkyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfonylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, aminoalkyl, alkoxyalky,  $CH_2$ heterocycloalkyl $COOR_{10}$  heterocycloalkyl $COOR_{10}$ , -COOH,  $-COR_5$ ,  $-C(O)OR_5$ ,  $CONHR_5$ ,  $-C(O)C(O)OR_5$ ,  $COONHR_5$ ,  $COON(R_5)OR_5$ ,  $COCON(R_5)OR_5$ , COCON(R

each R₅ is independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl and acyl each of which may be optionally substituted;

R<sub>10</sub> is selected from H, alkyl, acyl and aryl.

- 35. A pharmaceutical composition including a compound according to any one of claims 1 to 34 and a pharmaceutically acceptable diluent, excipient or carrier.
- 15 36. Use of a compound according to any one of claims 1 to 34 in the preparation of a medicament for the treatment of a disorder caused by, associated with or accompanied by disruptions of cell proliferation and/or angiogenesis.
  - 37. A use according to claim 36 wherein the disorder is a proliferative disorder.

38. A use according to claim 36 wherein the proliferative disorder is cancer.

- 39. Use of a compound according to claim 38 wherein the cancer is selected from breast cancer, lung cancer, ovarian cancer, prostate cancer, head and neck cancer, renal cancer, gastric cancer, colon cancer, pancreatic cancer and brain cancer.
- 40. A method of treatment of a disorder caused by, associated with or accompanied by disruptions of cell proliferation and/or angiogenesis in a patient the method including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.
- 41. A method according to claim 40 wherein the disorder is a proliferative disorder.
- 42. A method according to claim 40 wherein the disorder is cancer.

PCT/SG2004/000354

10

15

20

- 43. A method according to claim 42 wherein the cancer is selected from breast cancer, lung cancer, ovarian cancer, prostate cancer, head and neck cancer, renal cancer, gastric cancer, colon cancer, pancreatic cancer and brain cancer.
- 5 44. Use of a compound according to any one of claims 1 to 34 to modify deacetylase activity.
  - 45. A use according to claim 44 wherein the deacetylase activity is histone deacetylase activity.
  - 46. A use according to claim 44 wherein the deacetylase activity is class I histone deacetylase activity.
  - 47. A use according to claim 45 or 46 wherein the histone deacetylase is HDAC1.
  - 48. A use according to claim 45 or 46 wherein the histone deacetylase is HDAC8.
  - 49. A method of modifying deacetylase activity including contacting the deacetylase with a compound according to any one of claims 1 to 34.
  - 50. A method according to claim 49 wherein the deacetylase activity is histone deacetylase activity.
- 51. A method according to claim 49 wherein the deacetylase activity is class I histone deacetylase activity.
  - 52. A method according to claim 50 or 51 wherein the histone deacetylase is HDAC1.
  - 53. A method according to claim 50 or 51 wherein the histone deacetylase is HDAC8.
    - 54. A method of treatment of a disorder that can be treated by the inhibition of deacetylase activity in a patient including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.
- 55. A method according to claim 54 wherein the deacetylase activity is histone deacetylase activity.

130

- 56. A method of treatment of a disorder that is mediated by histone deacetylase activity in a patient including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.
- 57. 5 A method according to any one of claims 54 to 56 wherein the disorder is selected from the group consisting of proliferative disorders (e.g. cancer); Neurodegenerative diseases including Huntington's Disease, Polyglutamine diseases, Parkinson's Disease. Alzheimer's Disease, Seizures, Striatonigral degeneration, Progressive supranuclear palsy, Torsion dystonia, Spasmodic torticollis and dyskinesis, Familial tremor, Gilles de la Tourette syndrome, Diffuse Lewy body disease, Progressive supranuclear palsy, Pick's 10 disease, Intracerebral haemorrhage, Primary lateral sclerosis, Spinal muscular atrophy, Amyotrophic lateral sclerosis, Hypertrophic interstitial polyneuropathy, Retinitis pigmentosa, Hereditary optic atrophy, Hereditary spastic paraplegia, Progressive ataxia and Shy-Drager syndrome; Metabolic diseases including Type 2 diabetes; Degenerative 15 Diseases of the Eye including Glaucoma, Age-related macular degeneration. Rubeotic glaucoma, Intersitital keratitis, diabetic retinopathy; Inflammatory diseases and/or Immune system disorders including Rheumatoid Arthritis (RA), Osteoarthritis, Juvenile chronic arthritis, Graft versus Host disease, Psoriasis, Asthma, Spondyloarthropathy, Crohn's Disease, Inflammatory bowel disease, Colitis Ulcerosa, Alcoholic hepatitis, Diabetes, Sjoegrens's syndrome, Multiple Sclerosis, Ankylosing spondylitis, Membranous 20 glomerulopathy, Discogenic pain, Systemic Lupus Erythematosus; Disease involving angiogenesis including cancer, psoriasis, rheumatoid arthritis; Psychological disorders including bipolar disease, schizophrenia, mania, depression and dementia; Cardiovascular Diseases including Heart failure, restenosis and arteriosclerosis; Fibrotic 25 diseases including liver fibrosis, cystic fibrosis and angiofibroma; Infectious diseases including Fungal infections, such as Candida Albicans, Bacterial infections, Viral infections, such as Herpes Simplex, Protozoal infections, such as Malaria, Leishmania infection, Trypanosoma brucei infection, Toxoplasmosis and coccidiosis and Haematopoietic disorders including thalassemia, anemia and sickle cell anemia.

- 58. A method for inhibiting cell proliferation including administration of an effective amount of a compound according to any one of claims 1 to 34.
- 59. A method of treatment of a neurodegenerative disorder in a patient including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.

131

- 60. A method according to claim 59 wherein the neurodegenerative disorder is Huntington's Disease.
- 61. A method of treatment of an inflammatory disease and/or immune system disorder in a patient including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.
- 62. A method according to claim 61 wherein the inflammatory disease and/or immune system disorder is rheumatoid arthritis.
- 63. A method according to claim 61 wherein the inflammatory disease and/or immune system disorder is systemic lupus erythematosus.
  - 64. A method of treatment of a proliferative disorder in patient including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.

65. A method of treatment of cancer in patient including administration of a therapeutically effective amount of a compound according to any one of claims 1 to 34 to the patient.

- 20 66. A method according to claim 65 wherein the cancer is a hematologic malignancy.
  - 67. A method according to claim 66 wherein the hematologic malignancy is selected from a group consisting of B-cell lymphoma, T-cell lymphoma and leukemia.
- 25 68. A method according to claim 65 wherein the cancer is a solid tumor.
  - 69. A method according to claim 67 wherein the solid tumor is selected from a group consisting of breast cancer, lung cancer, ovarian cancer, prostate cancer, head and neck cancer, renal cancer, gastric cancer, colon cancer, pancreatic cancer and brain cancer.
- 30 70. Use of a compound according to claims 1 to 34 in the manufacture of medicaments for the induction of apoptosis of tumor cells.
  - 71. Use of a compound according to any one of claims 1 to 34 in the preparation of a medicament for the treatment of cancer.

5

132

- 72. A use according to claim 72 wherein the cancer is a hematologic malignancy.
- 73. A use according to claim 72 wherein the hematologic malignancy is selected from the group consisting of B-cell lymphoma, T-cell lymphoma and leukemia.
- 74. A use according toc laim 71 wherein the cancer is a solid tumor.
- 75. A use according to claim 74 wherein the solid tumor is selected from a group consisting of breast cancer, lung cancer, ovarian cancer, prostate cancer, head and neck cancer, renal cancer, gastric cancer, colon cancer, pancreatic cancer and brain cancer.
- 76. A method for the induction of apoptosis of tumor cells including contacting the tumor cells with an effective amount of a compund according to any one of claims 1 to 34.

10